



## The Food Allergy & Anaphylaxis Network (FAAN) A Fact Sheet for Emergency Responders

The Food Allergy & Anaphylaxis Network (FAAN) is a national nonprofit organization established to increase public awareness about food allergy and anaphylaxis, to provide education, and to advance research on behalf of all those affected by food allergies. For more information call FAAN at 1-800-929-4040 or visit the website at [www.foodallergy.org](http://www.foodallergy.org).

- Approximately 12 million Americans, or 4% of the general population, suffer from food allergy.
- Eight foods account for 90% of the allergic reactions: peanuts, tree nuts (walnuts, pecans, etc.), fish, shellfish, eggs, milk, soy, and wheat. Peanuts are the leading cause of severe allergic reactions, followed by shellfish, fish, tree nuts, and eggs.
- Individuals with food allergies and asthma appear to be at an increased risk for severe allergic reactions.
- Food allergy is the leading cause of anaphylaxis outside the hospital setting, accounting for an estimated 30,000 ER visits per year.
- It is estimated that as many as 100 to 200 people die each year from food allergic reactions. Approximately 50 people die from insect sting reactions.

### What happens in the body during a food-allergic reaction?

The immune system mistakenly believes that a harmless substance, in this case a food item, is harmful. In its attempt to protect the body, it creates specific IgE antibodies to that food. The next time the individual eats that food, the immune system releases massive amounts of chemicals and histamines in order to protect the body. These chemicals trigger a cascade of allergic symptoms that can affect the respiratory system, gastrointestinal tract, skin, or cardiovascular system.

# Medic Alert®



## **What are the common symptoms of a reaction?**

Symptoms range from mild to severe. The severe or life-threatening reactions are called "anaphylaxis". Symptoms can include tingling sensation in the mouth, swelling of the tongue and the throat, difficulty breathing, hives, vomiting, abdominal cramps, diarrhea, drop in blood pressure, and loss of consciousness. In rare cases, death occurs. Symptoms typically appear within minutes to two hours after the person has eaten the food to which he or she is allergic.

## **What is the best treatment for food allergy?**

Strict avoidance of the allergy-causing food is the only way to avoid a reaction. Reading ingredient labels for all foods is the key to maintaining control over the allergy. If a product doesn't have a label, allergic individuals should not eat that food. If the label contains unfamiliar terms, shoppers must call the manufacturer and ask for a definition or avoid eating that food.

## **What is the best treatment for a food allergy reaction?**

Epinephrine, also called "adrenaline", is the medication of choice for controlling a severe allergic reaction. It is available as an auto-injector by prescription as EpiPen®, EpiPen® Jr, and Twinject™.

MedicAlert Foundation, one of the largest nonprofit membership organizations in the world, is the nation's leading emergency medical information service and has helped save lives for more than 40 years. As part of our lifesaving mission, we provide training materials to emergency responders through our Professional Education Department. For additional copies of this and other educational materials, please e-mail [profed@medicalert.org](mailto:profed@medicalert.org) or visit our Web site at [www.medicalert.org](http://www.medicalert.org).

Disclaimer: The information provided herein is not intended as a substitute for professional and medical advice, diagnosis, or treatment. Always seek the advice of your physician or other qualified health provider with any questions you may have regarding a medical condition. Never disregard professional medical advice or delay in seeking it because of something you have read in this material. If you think you have a medical emergency, call your doctor or 911 immediately. MedicAlert Foundation does not recommend or endorse any specific tests, products or procedures, opinions, or other information that may be stated in the material. Reliance on any information provided within this material is solely at your own risk. MedicAlert Foundation will not be held responsible for any action taken by readers as a result of their interpretation of the material, which is intended for educational and information purposes only.