







Aqoonso oo ka Jawaab xasaasiyadda daran Anaphylaxis

Fal-celinta xasaasiyadda cuntada ee la tuhunsan yahay ama firfircoon

MID WALBA EE CALAAMADAHA DARAN SOO SOCDA

-  **SANBABADA:** Neefta oo ku qabata, hiinraag, qufac soo noqnoqda
-  **Wadnaha:** Maqaar guduudan ama midab buluug ah, suuxid darem, garaaca wadnaha oo daciif ah, dawakhaad
-  **DHUUNTA:** Dhuunta oo xanniban ama xabeeb leh, dhibaato xagga neefsashada ama liqidda
-  **AFKA:** Barar weyn ee carrabka ama dibnaha
-  **MAQAARKA:** Firiiric cuncun leh meelo badan ee jirka ah, casaan baahsan
-  **CALOOSHA:** Matag soo noqnoqda, shuban daran
-  **WAX KALE:** Dareenka in wax xun dhowaan dhici doono, walaac, jahwareer

AMA MID KA BADAN CALAAMADAHA FUDUD

-  **SANKA:** Cuncunka sanko ama diif, hindhiso
-  **AFKA:** Cuncunka afka
-  **MAQAARKA:** Firiiric yar, cuncun yar
-  **CALOOSHA:** Lalabo khafiif ah ama dareen raaxo la'aan

1

ISLA MARKIIBA SII EPINEPHRINE

2

 **Wac 911**

Codso ambalaas wata epinephrine.

Tixgeli Daawooyin Dheeraad ah

(Epinephrine ka dib):

- » Antihistamine
- » Daawada neefta (bronchodilator) haddii neef la qabo

Sida loo meeleeyo jirka

- » Qofka jiifi oo lugaha kor u qaad.
- » Haddii neefsashadu ku adag tahay ama ay matagayaan, ha fadhiistaan ama dhinac u jiifi.

Tallaabooyinka Xiga

- » Haddii calaamadaha aysan soo roonaan, ama calaamadaha soo noqdaan, qiyaaso kale ee epinephrine ah ayaa la siin karaa 5 daqiiqo ama ka badan ka dib goorta la siiyay qiyaasta ugu dambeysey.
- » U qaad qaybta gurmada ee Isbitaalka (ER) oo halkaa joog ugu yaraan 4 saacadood sababtoo ah calaamaduhu way soo laaban karaan. Isla markiiba ogeysii xiriiryadooda degdega ah.

Ha ku tiirsanin daawada antihistamines.

Markaad shaki gasho, qofka sii daawada epinephrine oo wac 911.

