

## Food Allergy Facts and Statistics for the U.S.

### What Is a Food Allergy?

- A food allergy is an adverse health effect resulting from a specific immune response that occurs reproducibly on exposure to a given food.<sup>1</sup> The health effect, called an allergic reaction, occurs because the immune system attacks food proteins that are normally harmless.
- Food-induced anaphylaxis is a serious allergic reaction that is sudden in onset and may cause death.<sup>1</sup>
- There is no cure for food allergy. Strict avoidance of food allergens and early recognition and management of allergic reactions to food are important measures to prevent serious health consequences.<sup>2</sup>

### To Which Foods Are People Allergic?

- More than 170 foods have been reported to cause reactions in the U.S.<sup>1</sup>
- Eight major food allergens – milk, egg, peanut, tree nuts, wheat, soy, fish and crustacean shellfish – are responsible for most of the serious food allergy reactions in the United States.<sup>1</sup>
- Milk, egg, and peanut account for the majority of allergic reactions in young children.<sup>1</sup>
- Peanut, tree nuts, and seafood (fish and crustacean shellfish) account for the majority of reactions in teenagers and adults.<sup>1</sup>
- Allergists consider sesame allergy to be an emerging concern. Sesame has caused severe reactions, including fatal anaphylaxis.<sup>3,4,5,6</sup>

### How Many People Have Food Allergies?

- Approximately 15 million people in the United States have food allergies.<sup>7,8,9</sup>
- Roughly 4 percent of people age 18 or older – more than 9 million adults – have food allergies.<sup>7,8</sup>
- Results from a 2009-2010 survey of more than 38,000 children indicate that 5.9 million children, or 8 percent, have food allergies.<sup>8,9</sup> That's 1 in 13 children, or roughly two in every classroom.
- In a 2007 survey of 9,500 children, 3.9 percent were reported to have a food or digestive allergy within the past year.<sup>10</sup>
- About 30 percent of children with food allergies have multiple food allergies (more than one food to which they're allergic).<sup>9</sup>

### Food Allergies Are on the Rise

- The Centers for Disease Control & Prevention report that between 1997-1999 and 2009-2011, food allergy prevalence among children increased by 50 percent.<sup>11</sup>
- In the United States, the prevalence of childhood peanut or tree nut allergy appears to have more than tripled between 1997 and 2008.<sup>12</sup>
- The prevalence of childhood food allergy has increased at a rate of 2.1 percent per decade among blacks, 1.2 percent per decade among Hispanics and 1 percent per decade among whites, according to a study of self-reported allergy.<sup>13</sup>

## **Food Allergy Is a Serious Public Health and Economic Issue**

- A food allergy is an impairment that limits a major life activity and may qualify an individual for protection under the Americans with Disabilities Act of 1990 (ADA) and Section 504 of the Rehabilitation Act of 1973.<sup>14</sup>
- Caring for children with food allergies costs U.S. families nearly \$25 billion annually.<sup>15</sup>

## **Food Allergy Reactions Are Serious and Can Be Life-Threatening.**

- Every three minutes, a food allergy reaction sends someone to the emergency room.<sup>16</sup>
- Each year in the U.S., 200,000 people require emergency medical care for allergic reactions to food.<sup>16</sup>
- Pediatric hospitalizations for food allergy tripled between the late 1990s and the mid-2000s. Between 2004 and 2006, an average of 9,500 children received in-patient hospital care for food allergies each year.<sup>10</sup>
- About 40 percent of children with food allergies have experienced a severe allergic reaction such as anaphylaxis.<sup>9</sup>

## **Serious Allergic Reactions (Anaphylaxis) Require Immediate Treatment**

- Prompt injection of epinephrine (adrenaline) within minutes of the onset of anaphylaxis symptoms is crucial to successfully treating an anaphylactic reaction. A self-injectable epinephrine device is available by prescription.<sup>17</sup>
- Not recognizing the severity of an anaphylactic reaction and treating promptly (i.e., within minutes) with epinephrine is a risk factor for fatalities.<sup>18,19,20</sup>
- More than one dose of epinephrine may be required.<sup>21</sup>
- It is possible to have anaphylaxis without any skin symptoms, such as rash or hives.<sup>22</sup>
- Symptoms of anaphylaxis may recur after initially subsiding (known as a biphasic reaction). Experts recommend an observation period of 4 to 6 hours in the emergency room to monitor that the reaction has been resolved.<sup>21,23</sup>

## **Food Allergy Impacts Quality of Life**

- About 1 in 3 children with food allergy reports being bullied as a result. Among children with allergies to more than two foods, over half report being bullied due to food allergy.<sup>24</sup>
- Compared to children who do not have a medical condition, children with food allergy are twice as likely to be bullied.<sup>25</sup>
- More than one-quarter of parents surveyed during food allergy appointments report that their children do not participate in camp or sleepovers because of food allergy. More than 15 percent do not go to restaurants, and more than 10 percent avoid child care settings or playdates at friends' houses. Ten percent home-school their children to prevent food allergen exposure.<sup>26</sup>
- Among parents of young children in the first year after food allergy diagnosis, most avoid restaurants and about half restrict social activities or travel.<sup>25</sup>
- Mothers of food-allergic children under age five have significantly higher blood-pressure measurements and report significantly greater levels of psychosocial stress than mothers whose preschool-aged children do not have food allergies.<sup>27</sup>

### Who is at Highest Risk for Developing Food Allergy?

- Compared to white children, Asian and black children are more likely to report food allergy symptoms but are less likely to be diagnosed with food allergy by a physician.<sup>9</sup>
- Children from rural communities are less likely to have food allergies than children from urban centers.<sup>28</sup>
- Children from households earning less than \$50,000 per year are less likely to report food allergies than are children from households earning more than \$50,000 per year.<sup>9</sup>
- Among inner-city children with a family history of hay fever, eczema or asthma, 1 preschool-aged child in 10 is allergic to eggs, milk, or peanuts.<sup>29</sup>
- Compared to children without food allergy, children with food allergy are more than twice as likely to have asthma and more than three times as likely to have respiratory allergy or eczema.<sup>10</sup>
- Food allergies may trigger or be linked to eosinophilic gastrointestinal diseases.<sup>30</sup>
- While most common in childhood, a food allergy can develop at any time. Research suggests that at least 15 percent of food allergies are first diagnosed in adulthood.<sup>31</sup>

### Who Is at Highest Risk for Fatal Anaphylaxis?

- Although a severe or fatal reaction can happen at any age, teenagers and young adults with food allergies are at the highest risk of fatal food-induced anaphylaxis.<sup>18,19,20</sup>
- Individuals with food allergies who also have asthma may be at increased risk for severe or fatal food allergy reactions.<sup>18,20</sup>

### Under What Circumstances Do Reactions Occur?

- Food allergy reactions typically involve foods that are believed to be safe. Allergic reactions can result from mislabeling or cross-contact during food preparation.<sup>19,32,33,34</sup>
- Limited skin contact with peanut butter or inhaling peanut butter from a short distance is unlikely to elicit a significant allergic reaction. These results cannot be generalized to more extensive contact or to other forms of peanut.<sup>35,36</sup> *Note: Limited contact with peanut butter presents a greater risk to young children, who frequently put their hands in their mouths.*
- Food proteins released into the air in vapor or steam from cooked foods can potentially cause allergic reactions. Reactions from vapor or steam can resemble reactions to inhaled allergens that cause hay fever or asthma symptoms, such as pollen or animal dander.<sup>37,38</sup>

### Where Do Reactions Occur?

- Reports suggest that the majority of fatal food allergy reactions are triggered by food consumed outside the home.<sup>18,19,20</sup>
- One study looking at peanut and tree nut allergy reactions in restaurants and other food establishments found that reactions were frequently attributed to desserts, that Asian restaurants and take-out dessert stores (bakeries, ice cream shops) were common sources of foods that triggered reactions, and that the food establishment was often not properly notified of a food allergy by the customer with the allergy.<sup>39</sup>
- Research on self-reported reactions occurring on commercial airlines indicates that peanut and tree nut reactions on planes have resulted from ingestion, contact and inhalation. Ingestion of an allergen remains the main concern for severe reactions.<sup>40,41,42</sup>

## Are Food Allergy Reactions Common at School?

- More than 15 percent of school-aged children with food allergies have had a reaction in school.<sup>43,44</sup>
- In a 2013-2014 survey of schools participating in a program to provide undesignated (stock) epinephrine for emergency use, over 600 schools – more than 10 percent – reported at least one case of anaphylaxis.<sup>45</sup>
- Approximately 20-25 percent of epinephrine administrations in schools involve individuals whose allergy was unknown at the time of the reaction.<sup>46</sup>
- In one large school district during the 2012-2013 school year, more than half of the 38 individuals who were treated with district-supplied emergency epinephrine were experiencing their first severe reaction.<sup>47</sup>
- Food allergy reactions can happen in multiple locations throughout the school, and are not limited to the cafeteria. Care must be exercised during bake sales, classroom parties and opportunities for snacking.<sup>20,46</sup>

## Avoiding Allergens Requires Careful Reading of Labels and Stringent Cleaning Procedures.

- Even trace amounts of a food allergen can cause a reaction.<sup>48,49,50,51,52,53</sup>
- Some studies have shown that most individuals with peanut and soy allergies can safely eat highly refined oils made from these ingredients. However, cold-pressed, expeller-pressed, or extruded oils should be avoided. Talk to your doctor about oils made from ingredients to which you are allergic.<sup>54,55,56,57,58,48</sup>
- According to the Food Allergen Labeling and Consumer Protection Act (FALCPA) the major eight allergens must be declared in simple terms, either in the ingredient list or via a separate allergen statement. However, FALCPA does not regulate the use of advisory/precautionary labeling (e.g., “may contain,” “made in a facility that also processes”).<sup>59</sup>
- Advisory/precautionary labeling is voluntary. The terms do not reflect specific risks, and random product testing has found allergen levels ranging from undetectable to amounts that can cause allergic reactions.<sup>1,32</sup>
- A study showed that peanut protein was detected in 7.3 percent of products bearing advisory/precautionary labeling for peanut.<sup>33</sup>
- A study showed that peanut can be cleaned from the hands of adults by using running water and soap or commercial wipes, but not by applying antibacterial gels. In addition, peanut was easily removed from surfaces by using common household cleaning sprays or sanitizing wipes but not by wiping with dishwashing liquid.<sup>60</sup>

## Can Food Allergies Be Outgrown?

- Although allergies to milk, egg, wheat and soy often resolve in childhood, research suggests that children may outgrow at least some of these food sensitivities more slowly than was found in previous decades, with many children still allergic beyond age 5.<sup>1</sup>
- Allergies to peanuts, tree nuts and shellfish are generally lifelong.<sup>1</sup>

## Food Allergy Prevention and Treatment

- **Learning Early About Peanut Allergy (LEAP)** provided evidence that the age at which a child first eats peanut and the frequency of peanut in the diet can influence whether the child develops an allergy to peanut. LEAP findings demonstrate that early, sustained consumption

of peanut products is associated with a substantial and significant decrease in the likelihood of developing peanut allergy.<sup>61</sup>

- In 2017, findings from LEAP and related studies led to the release of new guidelines for introduction of peanut.<sup>62</sup>
- A follow-up to the LEAP trial, **Persistence of Oral Tolerance to Peanut (LEAP-On)**, showed that decreased peanut allergy risk among children who consumed peanut throughout early childhood persists even after the children avoid peanut from ages 5 to 6.<sup>63</sup>
- Food allergy therapies are under study in clinical trials, but none has been proven yet for general use\*. Some therapies under investigation include:
  - **Oral immunotherapy (OIT)** – To raise the threshold dose at which food allergy reactions occur, progressively greater amounts of allergen are eaten under medical supervision until a maintenance dose is reached. Success in experimental OIT studies, resulting in a significant increase in the tolerated threshold dose of the problem food, can vary from 30 to 60%.<sup>64</sup> Successful OIT can result in desensitization, that is, the ability to eat a significant amount of the problem food without a reaction. Desensitization can be lost if the problem food is not consumed on an ongoing basis. Side effects can be severe, including anaphylaxis and eosinophilic esophagitis.<sup>65</sup>
  - **Sublingual immunotherapy (SLIT)** – Food protein in liquid form is held under the tongue for a time before being swallowed. The desensitization achieved with SLIT is generally less robust than desensitization achieved with OIT, but SLIT is less likely to cause serious allergic reactions.<sup>66</sup>
  - **Epicutaneous immunotherapy (EPIT, or skin patch)** – Research is being conducted to evaluate the safety and efficacy of dermal patches that deliver allergens through the skin.<sup>67</sup>

\*FARE notes that oral and sublingual immunotherapy for food allergy are being conducted in private practices. To find FARE's position on OIT in private practice, please visit [www.foodallergy.org/press-room/fare-statements/021816](http://www.foodallergy.org/press-room/fare-statements/021816).

---

<sup>1</sup> NIAID-Sponsored Expert Panel. Guidelines for the diagnosis and management of food allergy in the United States: Report of the NIAID-sponsored expert panel. *J Allergy Clin Immunol*. 2010; 126(6):S1-58

<sup>2</sup> U.S. Food and Drug Administration, Center for Food Safety and Applied Nutrition. Food allergies: What you need to know. 2008. Retrieved from <http://www.cfsan.fda.gov/~dms/ffa1rgn.html>

<sup>3</sup> Ben-Shoshan M, Harrington DW, Soller L, Fragapane J, Joseph L, St Pierre Y, Godefroy SB, Elliott SJ, Clarke AE. A population-based study on peanut, tree nut, fish, shellfish, and sesame allergy prevalence in Canada. *J Allergy Clin Immunol*. 2010; 125(6):1327-1335.

<sup>4</sup> Morisset M, Moneret-Vautrin DA, Kanny G, Guénard L, Beaudouin E, Flabbée J, Hatahet R. Thresholds of clinical reactivity to milk, egg, peanut and sesame in immunoglobulin E-dependent allergies: evaluation by double-blind or single-blind placebo-controlled oral challenges. *Clin Exp Allergy*. 2003; 33(8):1046-1051.

<sup>5</sup> Gangur V, Kelly C, Navuluri L. Sesame allergy: a growing food allergy of global proportions? *Ann Allergy Asthma Immunol*. 2005; 95(1):4-11.

<sup>6</sup> Patel A, Bahna SL. Hypersensitivities to sesame and other common edible seeds. *Allergy*. 2016 e-pub ahead of publication. doi: 10.1111/all.12962.

- <sup>7</sup> National Institute of Allergy and Infectious Diseases, National Institutes of Health. Report of the NIH Expert Panel on Food Allergy Research. 2006. Retrieved from [www.niaid.nih.gov/topics/foodallergy/research/pages/reportfoodallergy.aspx](http://www.niaid.nih.gov/topics/foodallergy/research/pages/reportfoodallergy.aspx)
- <sup>8</sup> United States Census Bureau Quick Facts (2015 estimates).
- <sup>9</sup> Gupta RS, Springston MR, Warrier BS, Rajesh K, Pongracic J, Holl JL. The prevalence, severity, and distribution of childhood food allergy in the United States. *Pediatrics* 2011; 128(1):e9-17.
- <sup>10</sup> Branum A, Lukacs S. Food allergy among U.S. children: Trends in prevalence and hospitalizations. NCHS data brief, no 10. Hyattsville, MD: National Center for Health Statistics. 2008. Retrieved from [www.cdc.gov/nchs/data/databriefs/db10.htm](http://www.cdc.gov/nchs/data/databriefs/db10.htm)
- <sup>11</sup> Jackson KD, Howie LD, Akinbami LJ. Trends in allergic conditions among children: United States, 1997-2011. NCHS data brief, no 121. Hyattsville, MD: National Center for Health Statistics. 2013. Retrieved from <http://www.cdc.gov/nchs/products/databriefs/db121.htm>.
- <sup>12</sup> Sicherer SH, Muñoz-Furlong A, Godbold JH, Sampson HA. US prevalence of self-reported peanut, tree nut, and sesame allergy: 11-year follow-up. *J Allergy Clin Immunol.* 2010; 125(6):1322–1326.
- <sup>13</sup> Keet CA, Savage JH, Seopaul S, Peng RD, Wood RA, Matsui EC. Temporal trends and racial/ethnic disparity in self-reported pediatric food allergy in the United States. *Ann Allergy Asthma Immunol.* 2014 Mar; 112(3):222-229.
- <sup>14</sup> U.S. Department of Education, Office for Civil Rights. Questions and Answers on the ADA Amendments Act of 2008 for Students with Disabilities Attending Public Elementary and Secondary Schools. <http://www2.ed.gov/about/offices/list/ocr/docs/dcl-504faq-201109.html>, retrieved December 28, 2015.
- <sup>15</sup> Gupta R, Holdford D, Bilaver L, Dyer A, Holl JL, Meltzer D. The economic impact of childhood food allergy in the United States. *JAMA Pediatr.* 2013 Nov; 167(11):1026-31.
- <sup>16</sup> Clark S, Espinola J, Rudders SA, Banerji, A, Camargo CA. Frequency of US emergency department visits for food-related acute allergic reactions. *J Allergy Clin Immunol.* 2011; 127(3):682-683.
- <sup>17</sup> American Academy of Allergy, Asthma and Immunology, and American College of Allergy, Asthma and Immunology. Joint Task Force on Practice Parameters; Joint Council of Allergy, Asthma and Immunology. The diagnosis and management of anaphylaxis: an updated practice parameter. *J Allergy Clin Immunol.* 2005; 115:S483-523.
- <sup>18</sup> Bock SA, Muñoz-Furlong A, Sampson HA. Further fatalities caused by anaphylactic reactions to food, 2001–2006. *J Allergy Clin Immunol.* 2007; 119(4):1016-1018.
- <sup>19</sup> Bock SA, Muñoz-Furlong A, Sampson HA. Fatalities due to anaphylactic reactions to foods. *J Allergy Clin Immunol.* 2001; 107(1):191-193.
- <sup>20</sup> Sampson HA, Mendelson L, Rosen J. Fatal and near-fatal anaphylactic reactions to food in children and adolescents. *N Engl J Med.* 1992; 327(6):380-384.
- <sup>21</sup> Korenblat P, Lundie MJ, Danker RE, Day JH. A retrospective study of epinephrine administration for anaphylaxis: how many doses are needed? *Allergy Asthma Proc.* 1999; 20:383-386.
- <sup>22</sup> Sampson HA. Anaphylaxis and Emergency Treatment. *Pediatrics* 2003; 111(S6):1601-1608.
- <sup>23</sup> Ellis AK, Day JH. Incidence and characteristics of biphasic anaphylaxis: a prospective evaluation of 103 patients. *Ann Allergy Asthma Immunol.* 2007 Jan; 98(1) 64-69.
- <sup>24</sup> Shemesh E, Annunziato RA, Ambrose MA, Ravid NL, Mullarkey C, Rubes M, Chuang K, Sicherer M, Sicherer S. Child and parental reports of bullying in a consecutive sample of children with food allergy. *Pediatrics* 2013; 131:e10-e17.

- <sup>25</sup> Herbert L, Shemesh E, Bender B. Clinical management of psychosocial concerns related to food allergy. *J Allergy Clin Immunol Pract*. 2016; 4(2):205-213.
- <sup>26</sup> Bollinger ME; Dahlquist LM, Mudd K; Sonntag C, Dillinger L, McKenna K. The impact of food allergy on the daily activities of children and their families. *Ann Allergy Asthma Immunol*. 2006; 96:415-421.
- <sup>27</sup> Walker SO, Mao G, Caruso D, Hong X, Pongracic JA, Wang X. Cardiovascular risk factors in parents of food-allergic children. *Medicine (Baltimore)*. 2016 Apr; 95(15): e3156.
- <sup>28</sup> Gupta RS, Springston EE, Smith B, Warriar MR, Pongracic J, Holl JL. Geographic variability of childhood food allergy in the United States. *Clin Pediatr (Phila)*. 2012; 51(9):856-861.
- <sup>29</sup> McGowan EC, Bloomberg GR, Gergen PJ, Visness CM, Jaffee KF, Sandel M, O'Connor G, Kattan M, Gern J, Wood RA. Influence of early-life exposures on food sensitization and food allergy in an inner-city birth cohort. *J Allergy Clin Immunol*. 2015; 135(1):171-178.
- <sup>30</sup> Liacouras CA, Furuta GT, Hirano I, Atkins D, Attwood SE, Bonis PA, Burks AW, Chehade M, Collins MH, Dellon ES, Dohil R, Falk GW, Gonsalves N, Gupta SK, Katzka DA, Lucendo AJ, Markowitz JE, Noel RJ, Odze RD, Putnam PE, Richter JE, Romero Y, Ruchelli E, Sampson HA, Schoepfer A, Shaheen NJ, Sicherer SH, Spechler S, Spergel JM, Straumann A, Wershil BK, Rothenberg ME, Aceves SS. Eosinophilic esophagitis: Updated consensus recommendations for children and adults. *J Allergy Clin Immunol*. 2011 Jul; 128(1):3-20.
- <sup>31</sup> Kamdar TA, Peterson S, Lau CH, Saltoun CA, Gupta RS, & Bryce PJ. Prevalence and characteristics of adult-onset food allergy. *J Allergy Clin Immunol Pract*. 2015; 3(1):114–115.e1.
- <sup>32</sup> Ford LS, Taylor SL, Pacenza R, Niemann LM, Lambrecht DM, Sicherer SH. Food allergen advisory labeling and product contamination with egg, milk, and peanut. *J Allergy Clin Immunol*. 2010; 126(2):384-385.
- <sup>33</sup> Hefle SL, Furlong TJ, Niemann L, Lemon-Mule H, Sicherer S, and Taylor SL. Consumer attitudes and risks associated with packaged foods having advisory labeling regarding the presence of peanuts. *J Allergy Clin Immunol*. 2007; 120:171-176.
- <sup>34</sup> Taylor SL, Baumert JL. Cross-contamination of foods and implications for food allergic patients. *Curr Allergy Asthma Rep*. 2010 Jul; 10(4):265-70.
- <sup>35</sup> Simonte SJ, Sonhui M, Shideh M, Sicherer S. Relevance of casual contact with peanut butter in children with peanut allergy. *J Allergy Clin Immunol*, 2003; (112):180-182.
- <sup>36</sup> Wainstein BK, Kashef S, Ziegler M, Jelley D, Ziegler JB. Frequency and significance of immediate contact reactions to peanut in peanut-sensitive children. *Clin Exp Allergy*. 2007; 37(6):839–845.
- <sup>37</sup> Crespo JF, Pascual C, Dominguez C, Ojeda I, Munoz FM, Esteban MM. Allergic reactions associated with airborne fish particles in IgE-mediated fish hypersensitive patients. *Allergy*. 1995; 50(3):257-61.
- <sup>38</sup> Roberts G, Golder N, Lack G. Bronchial challenges with aerosolized food in asthmatic, food-allergic children. *Allergy*. 2002; 57:713-7.
- <sup>39</sup> Furlong TJ, DeSimone J, Sicherer SH. Peanut and tree nut allergic reactions in restaurants and other food establishments. *J Allergy Clin Immunol*. 2001; 108:867-870.
- <sup>40</sup> Sicherer SH, Furlong TJ, DeSimone J, Sampson HA. Self-reported allergic reactions to peanut on commercial airliners. *J Allergy Clin Immunol*. 1999; 103(103):186-189.
- <sup>41</sup> Comstock SS, DeMera R, Vega L, Boren EJ, Deanne S, Haapanen LA, Teuber SS. Allergic reactions to peanuts, tree nuts, and seeds aboard commercial airliners. *Ann Allergy Asthma Immunol*. 2008; 101:51-56.
- <sup>42</sup> Greenhawt MJ, McMorris MS, Furlong TJ. Self-reported allergic reactions to peanut and tree nuts on commercial airlines. *J Allergy Clin Immunol*. 2009; 124(3):598-599.

- 43 Nowak-Wegrzyn A, Conover-Walker MK, Wood RA. Food-allergic reactions in schools and preschools. *Arch Pediatr Adolesc Med.* 2001; 155(7):790-795.
- 44 Sicherer SH, Furlong TJ, DeSimone J, Sampson HA. The US peanut and tree nut allergy registry: characteristics of reactions in schools and day care. *J Pediatr.* 2011; 138(4): 560-565.
- 45 White MV, Hogue SL, Bennett ME, Goss D, Millar K, Hollis K, Siegel PH, Wolf RA, Wooddell MJ, Silvia S. EpiPen4Schools pilot survey: Occurrence of anaphylaxis, triggers, and epinephrine administration in a U.S. school setting. *Allergy Asthma Proc.* 2015 Jul-Aug;36(4):306-312.
- 46 McIntyre CL, Sheetz AH, Carroll CR, Young MC. Administration of epinephrine for life-threatening allergic reactions in school settings. *Pediatrics* 2005; 116(5):1134-1140.
- 47 DeSantiago-Cardenas L, Rivkina V, Whyte SA, Harvey-Gintoft BC, Bunning BJ, Gupta RS. Emergency Epinephrine Use for Food Allergy Reactions in Chicago Public Schools. *Amer J Prev Med.* 2015; 48(2): 170-173.
- 48 Hefle SL, Taylor SL. Allergenicity of edible oils. *Food Technol.* 1999; 53:62-70.
- 49 Laoprasert N, Wallen ND, Jones RT, Hefle SL, Taylor SL, Yunginger JW. Anaphylaxis in a milk-allergic child following ingestion of lemon sorbet containing trace quantities of milk. *J Food Prot.* 1998; 61:1522-1524.
- 50 Gern JE, Yang E, Evrard HM, Sampson HA. Allergic reactions to milk-contaminated nondairy products. *N Engl J Med.* 1991; 324:976-979.
- 51 Yunginger JW, Gauerke MB, Jones RT, Dahlberg MJE, Ackerman SJ. Use of radioimmunoassay to determine the nature, quantity and source of allergenic contamination of sunflower butter. *J Food Prot.* 1983; 46:625-628.
- 52 Jones R, Squillace D, Yunginger J. Anaphylaxis in a milk-allergic child after ingestion of milk contaminated kosher-pareve-labeled "dairy-free" dessert. *Ann Allergy.* 1992; 68:223-227.
- 53 Hourihane J, Kilbrun S, Nordlee J, et al. An evaluation of the sensitivity of subjects with peanut allergy to very low doses of peanut: a randomized, double-blind, placebo-controlled food challenge study. *J Allergy Clin Immunol.* 1997; 100:596-600.
- 54 Bush RK, Taylor SL, Nordlee JA, Busse WW. Soybean oil is not allergenic to soybean-sensitive individuals. *J Allergy Clin Immunol.* 1985; 76:242-245.
- 55 Taylor SL, Busse WW, Sachs M, Parker JL, Yunginger JW. Peanut oil is not allergenic to peanut-sensitive individuals. *J Allergy Clin Immunol.* 1981; 68:372-375.
- 56 Hoffman DR, Collins-Williams C. Cold-pressed peanut oils may contain peanut allergen. *J Allergy Clin Immunol.* 1994; 93:801-802.
- 57 Keating MU, Jones RT, Worley NJ, Shively A, Yunginger JW. Immunoassay of peanut allergens in food-processing materials and finished foods. *J Allergy Clin Immunol.* 1990; 86:41-44.
- 58 Crevel RW, Kerkhoff MA, Koning MG. Allergenicity of refined vegetable oils. *Food Chem Toxicol.* 2000; 38(4):385-393.
- 59 U.S. Food and Drug Administration. Food allergen labeling and consumer protection act of 2004 (public law 108-282, title II). Retrieved from <http://www.fda.gov/food/labelingnutrition/FoodAllergensLabeling/GuidanceComplianceRegulatoryInformation/ucm106187.htm>
- 60 Perry TT, Conover-Walker MK, Pomes A, Chapman MD, Wood RA. Distribution of peanut allergen in the environment. *J Allergy Clin Immunol.* 2004; 113(5):973-976.
- 61 Du Toit G, Roberts G, Sayre PH, Bahnson HT, Radulovic S, Santos AF, Brough HA, Phippard D, Basting M, Feeney M, Turcanu V, Sever ML, Gomez Lorenzo M, Plaut M, Lack G for the LEAP Study



Team. Randomized trial of peanut consumption in infants at risk for peanut allergy. *New Engl J Med*. 2015; 372:803-813.

<sup>62</sup> Togias A, Cooper SF, Acebal ML, Assa'ad A, Baker JR, Beck LA, Block J, Byrd-Bredbenner C, Chan ES, Eichenfield LF, Fleischer DM, Fuchs GJ, Furuta GT, Greenhawt MJ, Gupta RS, Habich M, Jones SM, Keaton K, Muraro A, Plaut M, Rosenwasser LJ, Rotrosen D, Sampson HA, Schneider LC, Sicherer SH, Sidbury R, Spergel J, Stukus DR, Venter C, Boyce JA. Addendum guidelines for the prevention of peanut allergy in the United States: Report of the National Institute of Allergy and Infectious Diseases–sponsored expert panel. *J Allergy Clin Immunol*. 2017; 39(1):29-44.

<sup>63</sup> Du Toit G, Sayre PH, Roberts G, Sever ML, Lawson K, Bahnson HT, Brough HA, Santos AF, Harris KM, Radulovic S, Basting M, Turcanu V, Plaut M, Lack G for the Immune Tolerance Network LEAP-On Study Team. *N Engl J Med* 2016; 374:1435-1443.

<sup>64</sup> Lanser BJ, Wright BL, Orgel KA, Vickery BP, Fleischer DM. Current Options for the Treatment of Food Allergy. *Pediatr Clin North Am*. 2015; 62(6):1531-1549.

<sup>65</sup> Wood RA. Food allergen immunotherapy: Current status and prospects for the future. *J Allergy Clin Immunol*. 2016; 137(4): 973-982.

<sup>66</sup> Narisety SD, Frischmeyer-Guerrerio PA, Keet CA, Gorelik M, Schroeder J, Hamilton RG, Wood RA. A randomized, double-blind, placebo-controlled pilot study of sublingual versus oral immunotherapy for the treatment of peanut allergy. *J Allergy Clin Immunol*. 2015; 135(5):1275-1282.e6.

<sup>67</sup> Jones SM, Sicherer SH, Burks AW, Leung DYM, Lindblad RW, Dawson P, Henning AK, Berin MC, Chiang D, Vickery BP, Pesek RD, Cho CB, Davidson WF, Plaut M, Sampson HA, Wood RA for the Consortium of Food Allergy Research. Epicutaneous immunotherapy for the treatment of peanut allergy in children and young adults. *J. Allergy Clin Immunol*. 2016; doi.org/10.1016/j.jaci.2016.08.017.